

Group for Improvement of Intestinal Function and Treatment

(GIFT)

SickKids



School Readiness Package

This package will provide you with information to prepare for having a child with intestinal failure in the classroom. Please use this to guide an open discussion with caregivers about the particular needs for this child in your classroom and add this document to this child's individual education plan.

In this package you will find:

- · Links educational videos for staff and fellow students
- "About my child" pages to clarify the child's needs
- Copies of Emergency Procedure Cards

Videos

Adult Caregivers

This video provides educators and adults with general information about intestinal failure. Intestinal Failure: Video for Adult Caregivers

Schoolmates

This video is intended to help schoolmates to help them understand intestinal failure and what that looks like. We encourage teachers to show this video to students and parents in this child's classroom as directed and with the consent of caregivers. Intestinal Failure: Video for Classmates

Emergency Procedures

Central venous access device (CVAD) procedures include what to do if line breaks, is cut, or cracks, if the cap falls off, if central line is pulled out, if the dressing comes loose: <u>Emergency Central Venous Access Device (CVAD) Procedures</u>

Each child will carry with them a copy of instructions for emergency procedures in their backpack.

SickKids



About my child

These are the devices on my child's body







□ has a CVAD



 $\hfill\square$ is connected to TPN during school

Generally,

is disconnected from TPN at _____

 $\hfill\square$ is not connected to TPN during school



has a gastrostomy tube



 $\hfill\square$ is connected to feeds/vent bag during school

Generally,

has feeds from _____

is not connected to feeds/vent bag during school



 \Box has an ostomy



Generally, the ostomy is emptied every _____ hours

My child...

 \Box is independent in emptying ostomy pouch

 \Box needs reminders to empty ostomy pouch

requires assistance to empty ostomy pouch

If the ostomy pouch leaks or detaches please note this is NOT an emergency. Here are instructions on what to do...



receives medications during the school day

My child...

 \Box is independent in taking their medications

 \Box needs reminders to take their medications

requires a school nurse to take their medications

does NOT receive medications during the school day

My child...

☐ has dietary restrictions. Please only allow foods sent from home unless otherwise approved.

has fluid restrictions:
They are only allowed to drink ______

at school

They are NOT allowed to drink at school

□ does NOT have dietary restrictions

CENTRAL LINE EMERGENCY PLAN IF LINE BREAKS, IS CUT, OR CRACKS			
CALL Caregivers – Inform them that the child's central line is broken and they need to come as soon as possible. Caregivers: Main:	Take GREEN SCISSORS (Kelly clamps) from emergency kit and CLAMP child's central line between their body and the breakage.		
2 Have child remain calm. Tell the child their line is broken and it is important to stay still and lay down.	4 If unable to reach caregiver CALL 911 and inform operator that you have a medical emergency and that the child has a central line, it is broken and they need EMS support.		

CENTRAL LINE EMERGENCY PLAN				
IF LINE IS PULLED OUT				
CALL Caregivers – Inform them that the child's central line has been pulled out and they need to come as soon as possible. Caregivers: Main:	Apply pressure to both neck and the exit site. After 10 minutes, check if there is bleeding at exit site. If there is bleeding present, continue to apply pressure. Chest line is depicted above; for PICC line apply pressure to the neck and arm where exit site is located.			
2 Have child remain calm. Tell the child their line has come out and it is important to stay still and lay down.	4 If unable to reach caregiver CALL 911 – Inform operator that you have a medical emergency and that the child has a central line that has been pulled out and they need EMS support.			

CENTRAL LINE EMERGENCY PLAN			
IF CAP FALLS OFF			
	CVL Clamp CAP		
CALL Caregivers – Inform them that the cap to the child's central line has fallen off and to come as soon as possible. Caregivers: Main:Alternate:	3 Make sure CVL clamp is closed. Take an alcohol swab from emergency kit and scrub the end of the line where the CAP would generally sit for 15 seconds and allow to dry for 15 seconds.		
Have child remain calm. Tell them that the CAP has fallen off their line and it is important to sit still and wait for their caregiver to come get them.	4 Take new CAP, from emergency kit and screw to the end of the central line. As soon as possible, a trained caregiver should change cap using aseptic techniques.		

G-TUBE EMERGENCY CARE

IF G-TUBE COMES OUT OF BODY



CALL Caregivers – Inform them that the	
G-tube has fallen out and to come as	
soon as possible and to let them know	
when the tube was dislodged. (The tube	
MUST be replaced within 6 hours).	

Retrieve G-tube emergency kit for child;

which should include replacement foley-

type catheter, gauze, syringe and tape.

Size of G-tube :_____

Caregivers:

Main: ____

1

Alternate: ____

0

If you have been trained to replace tube, proceed and replace with appropriate size Foley-type catheter.

If you have NOT been trained apply gauze to the site to minimize leakage on skin and await caregiver or medical personnel.

4

If waiting for medical personnel/ caregiver to replace tube have child remain calm. Tell them that their G-tube came out and it is important to stay still and to wait for the caregiver to arrive.